



MICROCOPY RESOLUTION TEST CHART NATIONAL BUREAU OF STANDARDS-1963-A

CHARACTERIZATION OF BINARY COMPLETELY SYMMETRIC CONNECTED BLOCK DESIGNS WITH v = b BASED ON n = O(MOD v) EXPERIMENTAL UNITS

bу

A. Hedayat*
University of Illinois, Chicago
Department of Mathematics
Chicago, Illinois, U.S.A.

ELECTE JANO Z 1983

Approved for public release; distribution unlimited.



A CHARACTERIZATION OF BINARY COMPLETELY SYMMETRIC CONNECTED BLOCK DESIGNS WITH v = b BASED ON n = O(MOD v) EXPERIMENTAL UNITS

by

A. Hedayat Tuniversity of Illinois, Chicago Department of Mathematics Chicago, Illinois, U.S.A.

ABSTRACT

It is shown that among binary and connected block designs with and without equal block sizes and equal replications only a symmetric balanced incomplete block design produces a completely symmetric information matrix for the treatment effects whenever the number of blocks is equal to the number of treatments and the number of experimental units is a multiple of the number of treatments. This result contains a known result as a special case.

AND WORKENS INCOMES SCIENTIFIC RESEARCH (AFSC)

THE TO DTIC

THE TO DT

AMS 1980 Subject Classification: Primary 62KlO; secondary 05B05. Key words and phrases: Completely symmetric information matrix, C-matrix. connected, binary design, block design, incomplete block, BIB design.

Research is supported by Grant AFOSR 80-0170.



UNCLASSIFIED
SECURITY CLASSIFICATION OF THIS PAGE (When Date Entered)

REPORT DOCUMENTATION PAGE	READ INSTRUCTIONS BEFORE COMPLETING FORM			
1. REPORT NUMBER 2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER			
AFOSR-TR- 82-1075 AD-A123293	<u> </u>			
4. TITLE (and Subtitle)	5. TYPE OF REPORT & PERIOD COVERED			
A CHARACTERIZATION OF BINARY COMPLETELY SYMMETRIC CONNECTED BLOCK DESIGNS WITH v = b BASED ON	TECHNICAL			
$n = o(MOD \ v)$ EXPERIMENTAL UNITS	6. PERFORMING ORG. REPORT NUMBER			
7. Author(e) A.S. Hedayat	8. CONTRACT OR GRANT NUMBER(*)			
A.S. Hedayac	AFOSR-80-0170			
9. PERFORMING ORGANIZATION NAME AND ADDRESS	10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS			
Department of Mathematics	DEC4400D- 0004/AE			
University of Illinois at Chicago Chicago IL 60680	PE61102F; 2304/A5			
11. CONTROLLING OFFICE NAME AND ADDRESS	12. REPORT DATE			
Mathematical & Information Sciences Directorate	1982			
Air Force Office of Scientific Research Bolling AFB DC 20332	13. NUMBER OF PAGES			
14. MONITORING AGENCY NAME & ADDRESS(II different from Controlling Office)	15. SECURITY CLASS. (of this report)			
	UNCLASSIFIED			
	15a. DECLASSIFICATION/DOWNGRADING SCHEDULE			
	SCHEDULE ,			
16. DISTRIBUTION STATEMENT (of this Report)				
Approved for public release; distribution unlimited	•			
	/			
	·			
17. DISTRIBUTION ST. 4ENT (of the abetract entered in Block 20, if different from	m Report)			
18. SUPPLEMENTARY (CTES				
19. KEY WORDS (Continue on reverse side if necessary and identify by block number)	,			
Completely symmetric information matrix; C-matrix;				
block design; incomplete block; BIB design.				
20. ABSTRACT (Continue on reverse side if necessary and identify by block number)				
It is shown that among binary and connected block designs with and without equal				
block sizes and equal replications only a symmetric balanced incomplete block				
design produces a completely symmetric information effects whenever the number of blocks is equal to t				
the number of experimental units is a multiple of the number of treatments.				
This result contains a known result as a special case.				
· ·				



A CHARACTERIZATION OF BINARY COMPLETELY SYMMETRIC CONNECTED BLOCK DESIGNS WITH v = b BASED ON n = O(MOD v) EXPERIMENTAL UNITS

by

A. Hedayat University of Illinois, Chicago Department of Mathematics Chicago, Illinois, U.S.A.

1		ssion		•	
d.	NTIS Dyta	GRA&	:I	X	
] 1	ሆ••^ኍ	nounce Cleat	d ion		
<u> </u>				7	
. By					
Distribution/					
Availability Codes					
Di	st	Avail Spe	an cia:	d/or L	
1	2	1			
	7		1	•	

1. <u>INTRODUCTION</u>

Suppose v treatments are to be compared via n experimental units grouped into b blocks with k_j experimental units in block j. An arrangement of the v treatments into the n experimental units is called a <u>design</u>. Let D be the collection of all such designs and let r_{di} be the number of experimental units which design d assigns to treatment i. We associate the following matrix

$$C_{d} = R_{d} - N_{d}K^{-1}N_{d}$$

with design d, where R_d and K_d stand for diagonal matrices with diagonal elements $r_{d1}, r_{d2}, \ldots, r_{dv}$ and k_1, k_2, \ldots, k_b respectively and $N_d = (n_{dij})$, represents the $v \times b$ incidence matrix of the treatments versus blocks. Throughout the paper we shall adopt the following standard notation. A' and Tr A for the transpose and the trace of the matrix A respectively; I for the identity matrix and J for a matrix with all entries equal to one. The dimensions of the matrices will be clear from the context.

The matrix Cd plays an important role in the statistical analysis of the data collected under design d whenever the model of response is the usual homoscedastic linear additive model which specifies that the expectation of an observation on treatment i in block j is $\mu + \alpha_1 + \beta_1$ (general mean+treatment effect+block effect), where μ , $\alpha_{\underline{i}}$ and $\beta_{\underline{i}}$ are unknown constants, $1 \leq i \leq v$, $1 \le j \le b$ with the additional assumption that the n observations are uncorrelated with common variance σ^2 . Under this model the reduced normal equations for estimating the vector of treatment effects can be written as $C_d \hat{a} = Q_d$, where $Q_d = (T_{d1}, T_{d2}, \dots, T_{dd})$ T_{dv})' - $N_dK^{-1}(B_{d1}, B_{d2}, \dots, B_{db})$ '. Here T_{di} stands for the total of all observations obtained under treatment i and $B_{d,i}$ stands for the total of all observations produced in block j. We recall that if $q^{\dagger}\hat{\alpha}$ is an estimable linear contrast in treatment effects then under design d the BLU estimator of q'a is $q'\hat{a} = q'C_{\bar{a}}Q_{\bar{a}}$ with variance $q'\hat{\alpha} = q'C_{d}q\sigma^2$. Here A is a generalized inverse of A, that is, any matrix A as long as AA A = A. For this reason Cd is known as the information matrix of the estimable treatment contrasts.

As we see from the definition of $C_{\mathbf{d}}$ each d uniquely determines its associated $C_{\mathbf{d}}$ matrix. Fortunately, in general the converse is not true, that is, there are different designs which yield identical information matrices. Therefore, in such a situation the statistician has more than one design at his disposal for his client who insists on running a design with a certain form

for the information matrix of his design. One particular pattern which yields a great deal of optimality is complete symmetry by which we mean C_d to be of the form $a_1I + a_2J$. Kiefer (1975) proved that any design which yields a completely symmetric Ca matrix is universally optimal if $\operatorname{Tr} C_d$ is maximum in the class of competing designs. In this paper we shall prove that if b = v and $n = 0 \pmod{v}$ then C_d will be completely symmetric if and only if d is a symmetric BIB design. The sufficiency of the condition is obvious. To prove its necessity we need to recall some basic old results and prove some new ones which we shall do in Section 2. We would like to point out that our result contains the interesting result of Rao (1966), who proved that an equireplicate binary balanced design with b = v is a symmetric BIB design. Specifically, we shall show that to obtain the conclusion of Rao (1966) we can start with a much weaker assumption. Namely, rather than assuming that d is equally replicated we only have to assume that the number of available experimental units, n. is a multiple of v. Of course, somewhere in the process of the proof we will see that, under the given conditions, our assumptions and the assumptions of Rao (1966) are indeed equivalent. But a priori we did not have to start with the strong condition of equal replications. A result has to be established (Proposition 2.6) in order to enable us to weaken the a priori condition or to enlarge the class of competing designs.

2. CHARACTERIZATION

In the sequel for simplicity we shall drop the subscript d to the matrices and quantities associated with design d. In Propositions 2.1-2.6 we establish some results needed to prove our result stated in Theorem 2.1 which was given originally in Hedayat (1972). The stated results can be easily verified whenever the proof is omitted.

<u>Proposition 2.1.</u> If d is a completely symmetric connected design with

$$C = a_1 I + a_2 J$$

then,

$$a_1 = TrC/(v-1)$$
 and $a_2 = -TrC/v(v-1)$.

Proposition 2.2. If d is a binary completely symmetric connected design with

$$C = a_1 I + a_2 J$$

then.

$$a_1 = (n-b)/(v-1)$$
 and $a_2 = -(n-b)/v(v-1)$.

<u>Definition</u> 2.1. d is said to be a proper design if $k_j = k$.

Since for a proper design n = bk then we have

<u>Proposition</u> 2.3. If d is a binary, proper and completely symmetric connected design, then

$$C = \frac{b(k-1)}{v-1} I - \frac{b(k-1)}{v(v-1)} J.$$

Given that d is completely symmetric, then in some settings we can make useful inferences about R and K.

Proposition 2.4. If d is a binary, proper and completely symmetric connected design, then d is equireplicated.

<u>Proof</u>: $C_{ii} = r_i (1 - \frac{1}{k})$ since d is binary and proper. Also, $C_{ii} = (n-b)/v$ since d is completely symmetric. Thus, $r_i = n/v$.

<u>Proposition 2.5</u>. If d is a binary, proper, equireplicated and completely symmetric connected design, then d is a BIB design.

Proof: On one hand

$$C_{ii'} = -1(k) \sum_{j=1}^{b} n_{ij} n_{i'j}.$$

One the other hand

$$C_{11} = -(n-b)/v(v-1),$$

and hence

$$\lambda_{ii'} = \sum_{j=1}^{b} n_{ij} n_{i'j} = r(k-1)/(v-1).$$

<u>Proposition</u> 2.6. If d is a binary and completely symmetric connected design with b = v then d is equireplicated if $n \equiv 0 \pmod{v}$.

Proof: Since d is binary and completely symmetric, then

(1)
$$C_{11} = (n-b)/v = (n-v)/v = (n/v) - 1.$$

On the other hand.

(2)
$$C_{11} = r_1 - \sum_{j=1}^{b} \frac{n_{1j}}{k_j}$$
.

(1) and (2) imply that

$$r_i = r - 1 + \sum_{j=1}^{b} \frac{n_{i,j}}{k_j}$$
 where $r = n/v$.

Since r_i is an integer, this implies that $\sum_{j=1}^{b} \frac{n_{i,j}}{k_j}$ is an integer greater than or equal to one. However, we observe that

$$\sum_{i=1}^{v} \left\lceil \sum_{j=1}^{b} \frac{n_{i,j}}{k_{j}} \right\rceil = \sum_{j=1}^{b} \frac{1}{k_{j}} \sum_{i=1}^{v} \frac{1}{k_{j}} (k_{j}) = b = v,$$

that is, v (=b) terms ≥ 1 add up to v which implies that each term $\equiv 1$. Therefore, $r_i = r - 1 + 1 = r$; i = 1, 2, ..., v.

Theorem 2.1. If d is a binary and completely symmetric connected design with b = v and n = O(mod v), then d is a symmetric BIB design.

<u>Proof:</u> By Proposition 2.6, it is sufficient to prove that d is proper.

$$C = R - NK^{-1}N' = rI - NK^{-1}N', \qquad n = vr$$

$$= \frac{n-b}{v-1} I - \frac{n-b}{v(v-1)} J$$

$$= \frac{v(r-1)}{v-1} I - \frac{r-1}{v-1} J.$$

Thus

$$NK^{-1}N' = rI - \frac{v(r-1)}{v-1}I + \frac{r-1}{v-1}J$$
$$= \frac{v-r}{v-1}I + \frac{r-1}{v-1}J.$$

Since

$$\left\lceil \frac{\mathbf{v} - \mathbf{r}}{\mathbf{v} - \mathbf{1}} \mathbf{I} + \frac{\mathbf{r} - \mathbf{1}}{\mathbf{v} - \mathbf{1}} \mathbf{J} \right\rceil^{-1} = \frac{\mathbf{v} - \mathbf{1}}{\mathbf{v} - \mathbf{r}} \mathbf{I} - \frac{\mathbf{r} - \mathbf{1}}{\mathbf{r} (\mathbf{v} - \mathbf{r})} \mathbf{J}$$

we have

$$[NK^{-1}N']^{-1} = (N')^{-1}KN^{-1} = \frac{v-1}{v-r}I - \frac{r-1}{r(v-r)}J.$$

By premultiplying by N' and pastmultiplying by N the both sides of the above relation we obtain

$$K = \frac{v-1}{v-r} N'N - \frac{r-1}{r(v-r)} N'JN.$$

Note that J = 11' and $N'1 = (k_1, ..., k_b)$. Since K is a diagonal matrix with the 1th diagonal entry equal to k_i we thus observe that:

(3)
$$k_i = k_i(v-1)/(v-r) - k_i^2(r-1)/r(v-r),$$

 $i, j = 1, 2, ..., b; i \neq j.$

(4)
$$0 = \lambda_{ij}(v-1)/(v-r) - k_i k_j(r-1)/r(v-r)$$

Because $k_i > 0$, we conclude from (3) that $k_i = r$. Consequently, (4) implies that $\lambda_{i,j} = r(r-1)/(v-1)$. This completes the proof.

REFERENCES

Hedayat, A., (1972). Theory of Exact Designs, Department of Florida State University, Tallahassee, Florida, U.S.A.

Kiefer, J., (1975). Construction and optimality of generalized Youden designs, In A Survey of Statistical Designs and Linear Models (J. N. Srivastava, ed.), pp. 333-353, North-Holland. Amsterdam.

Rao, M.B., (1966). A note on equi-replicate balanced designs with b = v, <u>Bull</u>. <u>Calcutta Statist</u>. <u>Assoc</u>., <u>15</u>, 43-44.

REPORT DOCUMENTATION PAGE	READ INSTRUCTIONS BEFORE COMPLETING FORM			
1. REPORT NUMBER 2. GOVT ACCESSION	NO. 3. RECIPIENT'S CATALOG NUMBER			
A Characterization of binary completely symmetric connected block designs with v = b based on n = O(mod v) experiment units				
7. AUTHOR(a)	8. CONTRACT OR GRANT NUMBER(#)			
A. Hedayat 9. PERFORMING ORGANIZATION NAME AND ADDRESS	10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS			
Department of Mathematics, University of Illinois, Chicago Circle, Box 4348, Chicago, Illinois 60680				
11. CONTROLLING OFFICE NAME AND ADDRESS Air Force Office of Scientific Research	March 1982			
Bolling AFB, Washington, D.C. 20332	13. NUMBER OF PAGES			
14. MONITORING AGENCY NAME & ADDRESS(It different from Controlling Office	ce) 15. SECURITY CLASS. (of this report)			
	15a. DECLASSIFICATION/DOWNGRADING SCHEDULE			
16. DISTRIBUTION STATEMENT (of this Report)				
Approved for public release; distribution	unlimited.			
17. DISTRIBUTION STATEMENT (of the ebstract entered in Block 20, if different from Report)				
18. SUPPLEMENTARY NOTES				
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Completely symmetric information matrix, C-matrix, connected, binary design, block design, incomplete block, BIB design.				
It is shown that among binary and connected without equal block sizes and equal metric balanced incomplete block design properties information metric for the tree	ected block designs with replications only a symproduces a completely			

the number of blocks is equal to the number of treatments and the

number of experimental units is a multiple of the number of treatments. This result contains a known result as a special case.

